

Invariant quantization in one and two parameters on semisimple coadjoint orbits of simple Lie groups

J. Donin

Department of Mathematics, Bar-Ilan University,
52900 Ramat-Gan, Israel,

D. Gurevich,

ISTV, Université de Valenciennes,
59304 Valenciennes, France,

S. Shnider

Department of Mathematics, Bar-Ilan University,
52900 Ramat-Gan, Israel

Abstract

Let $\mathcal{A}$ be the function algebra on a semisimple orbit, M , in the coadjoint representation of a simple Lie group, G , with the Lie algebra $\mathfrak{g}$. We study one and two parameter quantizations of $\mathcal{A}$, $\mathcal{A}_h$, $\mathcal{A}_{t,h}$, such that the multiplication on the quantized algebra is invariant under action of the Drinfeld-Jimbo quantum group, $U_h(\mathfrak{g})$. In particular, the algebra $\mathcal{A}_{t,h}$ specializes at $h = 0$ to a $U(\mathfrak{g})$, or G , invariant quantization, $\mathcal{A}_{t,0}$.

We prove that the Poisson bracket corresponding to $\mathcal{A}_h$ must be the sum of the so-called r -matrix and an invariant bracket. We classify such brackets for all semisimple orbits, M , and show that they form a $\dim H^2(M)$ parameter family, then we construct their quantizations.

A two parameter (or double) quantization, $\mathcal{A}_{t,h}$, corresponds to a pair of compatible Poisson brackets: the first is as described above and the second is the Kirillov-Kostant-Souriau bracket on M . Not all semisimple orbits admit a compatible pair of Poisson brackets. We classify the semisimple orbits for which such pairs exist and construct the corresponding two parameter quantization of these pairs in some of the cases.

1 Introduction

Passing from classical mechanics to quantum mechanics involves replacing the commutative function algebra, $\mathcal{A}$, of classical observables on the appropriate phase space, M , with a noncommutative (deformed) algebra, $\mathcal{A}_t$, of quantum observables (see [BFFLS] where the deformation quantization scheme is developed). The algebra $\mathcal{A}$ is a Poisson algebra and the product in $\mathcal{A}_t$ is given by a power series in the formal parameter t with leading

term the original commutative product and with leading term in the commutator given by the Poisson bracket. If the classical system is invariant under a Lie group of symmetries G , the associated quantum system often retains the group of symmetries. In particular, the algebra $\mathcal{A}_t$ is often invariant under the action of G , or under the action of its universal enveloping algebra $U(\mathfrak{g})$.

Modern field-theoretical models and, in particular, the problem of incorporating gravity into a quantum field theory led to the requirement of deforming (quantizing) the group symmetry and the phase space themselves. This is one of the reasons for the interest in quantum groups. The quantum group, $U_h(\mathfrak{g})$, defined by Drinfeld and Jimbo is a deformation of $U(\mathfrak{g})$ as a Hopf algebra. The quantization of the phase space and its symmetry group corresponds to a $U_h(\mathfrak{g})$ invariant deformation of the algebra $\mathcal{A}_t$, which leads us to the problem of two parameter (or double) quantization, $\mathcal{A}_{t,h}$, of the function algebra for a $U(\mathfrak{g})$ invariant Poisson structure. In other words, the problem of two parameter quantization appears if we want to quantize the Poisson bracket in such a way that multiplication in the quantized algebra is invariant under the quantum group action.

In the present paper we investigate the problem of one and two parameter invariant quantizations of the Poisson function algebra on a semisimple orbit in the coadjoint representation of a simple Lie group.

In general, if M is a manifold on which a semisimple Lie group G acts, it is not true that there exists even a one parameter $U_h(\mathfrak{g})$ invariant quantization of the function algebra, $\mathcal{A}_h$. In [DGM] it is proven that such a quantization exists if $M = G/H$ and the Lie algebra of H contains a maximal nilpotent subalgebra. A $U_h(\mathfrak{g})$ invariant quantization of the algebra of holomorphic sections of a line bundle over a flag manifold is constructed by similar methods in [DG1]. In all these cases the Poisson bracket which is quantized is the r -matrix bracket, r_M , on M . This bracket is determined by the bivector field $(\rho \otimes \rho)(r)$ for $r \in \wedge^2 \mathfrak{g}$, the Drinfeld-Jimbo r -matrix (of the form (3.3)), and $\rho : \mathfrak{g} \rightarrow \text{Vect}(M)$ is the mapping defined by the action of G on M .

The Sklyanin-Drinfeld (SD) bracket on G is determined by the difference of two bivector fields on G which are the right and left invariant extensions of the Drinfeld-Jimbo classical r -matrix. For details see [LW]. In [DG2] it is shown that a one parameter quantization of the SD bracket exists for all semisimple orbits in coadjoint representation of G .

In the present paper we show that for the semisimple orbits, M , the SD bracket is only one from $\dim H^2(M)$ parameter family of Poisson brackets admitting $U_h(\mathfrak{g})$ invariant quantization.

For symmetric spaces M , the r -matrix bracket satisfies the Jacobi identity and hence defines a Poisson bracket. The existence of a one parameter $U_h(\mathfrak{g})$ invariant quantization with the r -matrix Poisson bracket is proven in [DS1]. That paper also proves that when M is a hermitian symmetric space there exists a two parameter $U_h(\mathfrak{g})$ invariant quantization. The hermitian symmetric spaces form a subclass of semisimple orbits in $\mathfrak{g}^*$. These orbits have a very interesting property: the one-sided invariant components of the Sklyanin-Drinfeld bracket being reduced on such an orbit become Poisson brackets separately. More precisely, one of these components being reduced on M is the r -matrix bracket and the other one becomes the Kirillov-Kostant-Souriau (KKS) bracket which is roughly speaking the restriction to the orbit of the Lie bracket in $\mathfrak{g}$. (see [KRR], [DG1]). These brackets are

obviously compatible (i.e. their Schouten bracket vanishes). So, we get a Poisson pencil which is the set of linear combinations of KKS and r -matrix Poisson brackets. The two parameter quantization, $\mathcal{A}_{t,h}$, for hermitian symmetric spaces constructed in [DS1] is just a quantization of such a pencil. In particular, $\mathcal{A}_{t,0}$ is a $U(\mathfrak{g})$ invariant quantization of the KKS bracket on M .

The orbits in $\mathfrak{g}^*$ on which the r -matrix bracket is Poisson have been classified in [GP]. In particular, the only such semisimple orbits are symmetric spaces.

It turns out that there exist two parameter $U_h(\mathfrak{g})$ invariant quantizations on some G -manifolds where the r -matrix bracket is not Poisson (does not satisfy the Jacobi identity) but one can add a G -invariant bracket to the r -matrix bracket and get a Poisson bracket. For example, in [Do] it is shown that for $\mathfrak{g} = \mathfrak{sl}(n)$ there is a two parameter $U_h(\mathfrak{g})$ invariant family $(S\mathfrak{g})_{t,h}$, where $S\mathfrak{g}$ is the symmetric algebra of $\mathfrak{g}$ which can be considered as the function (polynomial) algebra on $\mathfrak{g}^*$. It is proven that this family can be restricted to give a two parameter quantization on any semisimple orbit of maximal dimension (on which the r -matrix bracket is not Poisson).

In the present paper we prove the existence of a two parameter $U_h(\mathfrak{g})$ invariant quantization of the function algebra for some non-symmetric semisimple orbits in $\mathfrak{g}^*$. Similar to the case of symmetric orbits, the quasiclassical (infinitesimal) term of this quantization is a Poisson pencil generated by the KKS bracket and another Poisson bracket which must be the sum of r -matrix and an $U(\mathfrak{g})$ invariant brackets. In particular, we shall see that such pencils exist for all semisimple orbits in case $\mathfrak{g} = \mathfrak{sl}(n)$. Note, that for non-symmetric orbits both the r -matrix bracket and the SD Poisson bracket are not compatible with the KKS bracket. We give a complete classification of the orbits admitting such a Poisson pencil. Moreover, we classify all such pencils and construct the deformation quantization of some of them.

We now describe the content of the paper in more detail. Let G be a simple, connected, complex Lie group, $\mathfrak{g}$ its Lie algebra, The Drinfeld-Jimbo quantum group $U_h(\mathfrak{g})$ can be considered as algebra $U(\mathfrak{g})[[h]]$ with undeformed multiplication but deformed non-cocommutative comultiplication Δ_h .

Let M be a G -homogeneous complex manifold. It is easy to show that M is isomorphic to a semisimple orbit of G in the coadjoint representation $\mathfrak{g}^*$ if and only if the stabilizer, G_o , of a point $o \in M$ is a Levi subgroup in G (see Section 3 for definition). The symplectic structure on M is not unique, but each symplectic structure on M arises from an isomorphism of M with an orbit, O_λ , for some semisimple element $\lambda \in \mathfrak{g}^*$. On O_λ there is the Kirillov-Kostant-Souriau (KKS) Poisson bracket, v_λ , whose action on the restriction of linear functions to the orbit is given by the Lie bracket in $\mathfrak{g}$. Each symplectic structure on M is induced from the KKS bracket by an isomorphism of M onto a semisimple orbit (see Section 3).

The first problem we consider is that of quantizing the algebra $\mathcal{A}$ of polynomial (or holomorphic) functions on M , such that the quantized algebra $\mathcal{A}_h$ has a $U_h(\mathfrak{g})$ invariant multiplication

$$\mu_h = \sum_{i=0}^{\infty} h^i \mu_i,$$

where μ_0 is the initial multiplication of functions and the μ_i for $i \geq 1$ are bidifferential

operators. Invariance means that μ_h satisfies the property

$$\mu_h \Delta_h(x)(a \otimes b) = x \mu_h(a \otimes b) \quad \text{for } x \in U_h(\mathfrak{g}), \quad a, b \in \mathcal{A}.$$

The second problem we consider is the existence of a two parameter $U_h(\mathfrak{g})$ invariant quantization, $\mathcal{A}_{t,h}$, of $\mathcal{A}$ such that the one parameter family $\mathcal{A}_{t,0}$ is a $U(\mathfrak{g})$ invariant quantization of the KKS bracket on M .

The infinitesimal of the one parameter quantization $\mathcal{A}_h$ is a Poisson bracket, whereas the infinitesimal of the two parameter quantization is a pencil of two compatible Poisson brackets.

In Section 2 we recall some facts on the Drinfeld monoidal categories of quantum group representations, using in our construction of quantization. In addition, we show that the Poisson bracket, $p(a, b) = \mu_1(a, b) - \mu_1(b, a)$, corresponding to the $U_h(\mathfrak{g})$ invariant quantization $\mathcal{A}_h$ must be of a special form. Namely, let $r \in \wedge^2 \mathfrak{g}$ be the Drinfeld-Jimbo classical r -matrix. The Schouten bracket $\llbracket r, r \rrbracket \in \wedge^3 \mathfrak{g}$ is the invariant element φ , which is unique up to a factor. Denote by r_M the bracket on M determined by the bivector field $(\rho \otimes \rho)(r)$ where $\rho : \mathfrak{g} \rightarrow \text{Vect}(M)$ is the mapping defined by the action of G on M . We call r_M an r -matrix bracket. Put also $\varphi_M = \rho^{\otimes 3} \varphi$. Then, $p(a, b)$ has the form

$$p(a, b) = r_M(a, b) + f(a, b), \quad a, b \in \mathcal{A}, \quad (1.1)$$

where $f(a, b)$ is a $U(\mathfrak{g})$ invariant bracket with the Schouten bracket

$$\llbracket f, f \rrbracket = -\varphi_M. \quad (1.2)$$

Note that r -matrix bracket is compatible with any invariant bracket, i.e. $\llbracket f, r_M \rrbracket = 0$.

Similarly, the two parameter quantization $\mathcal{A}_{t,h}$ corresponds to a pair of compatible Poisson brackets, (p, v_λ) , where $p = r_M + f$ of the form (1.1) and v_λ is the KKS bracket. Since r_M is compatible with the invariant bracket v_λ , compatibility of p with v_λ is equivalent to condition

$$\llbracket f, v_\lambda \rrbracket = 0. \quad (1.3)$$

In Section 3 we give a classification of all invariant brackets f satisfying condition (1.2) for all M isomorphic to semisimple orbits. We show that such brackets form a $\dim H^2(M)$ parameter family.

In the same section we show that a semisimple orbit may not have any invariant brackets satisfying the conditions (1.2) and (1.3). We call an orbit a “good orbit”, if such a bracket exists and then give a classification of all good orbits. Namely, if $\mathfrak{g}$ is of type A_n , all semisimple orbits are good. All orbits which are symmetric spaces are good. In cases D_n , E_6 there are good orbits which are not symmetric spaces. Moreover, for the good orbits the brackets satisfying the conditions (1.2) and (1.3) form a one parameter family. In fact, the property of an orbit being good depends only on its structure as a homogeneous manifold, not on the symplectic structure. The dependence is on the Lie subalgebra of the stabilizer subgroup. So, if an orbit M is good, then any orbit isomorphic to M as a homogeneous manifold will be good.

In Section 4 we consider cohomologies of the complex of invariant polyvector fields on M with differential given by the Schouten bracket with the bivector f satisfying (1.2).

We show that for almost all f these cohomologies coincide with the usual de Rham cohomologies of the manifold M and then use this fact in Section 5 to prove the existence of an invariant quantization. In the proof we use methods of [DS1] and [DS2]. Using the same methods, we also construct the two parameter quantization for good orbits in cases D_n , E_6 and brackets satisfying (1.2) and (1.3). For the case A_n some additional arguments are required. (See [Do] where using another method the existence of two parameter quantization for maximal orbits is proven.)

In conclusion we make two remarks.

Remark 1. In the paper G is supposed to be a complex Lie group. However, one can consider the situation when G is a real simple Lie group with Lie algebra $\mathfrak{g}_{\mathbb{R}}$ and M is a semisimple orbit of G in $\mathfrak{g}_{\mathbb{R}}^*$. In this case we take $\mathcal{A} = C^\infty(M)$, the complex-valued smooth functions. Let $\mathfrak{g}$ be the complexification of $\mathfrak{g}_{\mathbb{R}}$. It is clear that Lie algebra $\mathfrak{g}$ and algebra $U(\mathfrak{g})$ act on $C^\infty(M)$. Since all our results are formulated in terms of $\mathfrak{g}$ action, they are valid in the real case as well (see also [DS1]).

Remark 2. The deformation quantization can be considered as the first step of a quantization procedure whose the second step is a representation of the quantized algebra, $\mathcal{A}_{t,h}$, as an operator algebra in a linear space. For some symmetric orbits in $sl(n)^*$ such a representation has been given in [DGR] and [DGK] but the method of [DGK] can be apparently extended to all symmetric orbits in $\mathfrak{g}^*$ for all simple Lie algebras $\mathfrak{g}$. These operator algebras have a deformed $U_h(\mathfrak{g})$ invariant trace which, however, is not symmetric.

2 Poisson brackets associated with $U_h(\mathfrak{g})$ invariant quantization

We recall some facts about the Drinfeld algebras and the monoidal categories determined by them. They will be used, in particular, in our construction of the quantization.

Let A be a commutative algebra with unit, B a unitary A -algebra. The category of representations of B in A -modules, i.e. the category of B -modules, will be a monoidal category if the algebra B is equipped with an algebra morphism, $\Delta : B \rightarrow B \otimes_A B$, called comultiplication, and an invertible element $\Phi \in B^{\otimes 3}$ such that Δ and Φ satisfy the conditions (see [Dr1])

$$(id \otimes \Delta)(\Delta(b)) \cdot \Phi = \Phi \cdot (\Delta \otimes id)(\Delta(b)), \quad b \in B, \quad (2.1)$$

$$(id^{\otimes 2} \otimes \Delta)(\Phi) \cdot (\Delta \otimes id^{\otimes 2})(\Phi) = (1 \otimes \Phi) \cdot (id \otimes \Delta \otimes id)(\Phi) \cdot (\Phi \otimes 1). \quad (2.2)$$

Define a tensor product functor for $\mathcal{C}$ the category of B modules, denoted $\otimes_{\mathcal{C}}$ or simply $\otimes$ when there can be no confusion, in the following way: given B -modules M, N , $M \otimes_{\mathcal{C}} N = M \otimes_A N$ as an A -module. The action of B is defined by

$$b(m \otimes n) = (\Delta b)(m \otimes n) = b_1 m \otimes b_2 n \quad \text{where } \Delta b = b_1 \otimes b_2,$$

using the Sweedler convention of an implicit summation over an index. The element $\Phi = \Phi_1 \otimes \Phi_2 \otimes \Phi_3$ defines the associativity constraint,

$$a_{M,N,P} : (M \otimes N) \otimes P \rightarrow M \otimes (N \otimes P), \quad a_{M,N,P}((m \otimes n) \otimes p) = \Phi_1 m \otimes (\Phi_2 n \otimes \Phi_3 p).$$

Again the summation in the expression for Φ is understood. By virtue of (2.1) Φ induces an isomorphism of B -modules, and by virtue of (2.2) the pentagon identity for monoidal categories holds. We call the triple (B, Δ, Φ) a Drinfeld algebra. The definition is somewhat non-standard in that we do not require the existence of an antipode. The category $\mathcal{C}$ of B -modules for B a Drinfeld algebra becomes a monoidal category. When it becomes necessary to be more explicit we shall denote $\mathcal{C}(B, \Delta, \Phi)$.

Let (B, Δ, Φ) be a Drinfeld algebra and $F \in B^{\otimes 2}$ an invertible element. Put

$$\tilde{\Delta}(b) = F\Delta(b)F^{-1}, \quad b \in B, \quad (2.3)$$

$$\tilde{\Phi} = (1 \otimes F) \cdot (id \otimes \Delta)(F) \cdot \Phi \cdot (\Delta \otimes id)(F^{-1}) \cdot (F \otimes 1)^{-1}. \quad (2.4)$$

Then $\tilde{\Delta}$ and $\tilde{\Phi}$ satisfy (2.1) and (2.2), therefore the triple $(B, \tilde{\Delta}, \tilde{\Phi})$ also becomes a Drinfeld algebra which has an equivalent monoidal category of modules, $\tilde{\mathcal{C}}(B, \tilde{\Delta}, \tilde{\Phi})$. Note that the equivalent categories $\mathcal{C}$ and $\tilde{\mathcal{C}}$ consist of the same objects as B -modules, and the tensor products of two objects are isomorphic as A -modules. The equivalence $\mathcal{C} \rightarrow \tilde{\mathcal{C}}$ is given by the pair (Id, F) , where $Id : \mathcal{C} \rightarrow \tilde{\mathcal{C}}$ is the identity functor of the categories (considered without the monoidal structures, but only as categories of B -modules), and $F : M \otimes_{\mathcal{C}} N \rightarrow M \otimes_{\tilde{\mathcal{C}}} N$ is defined by $m \otimes n \mapsto F_1 m \otimes F_2 n$ where $F_1 \otimes F_2 = F$.

Assume $\mathcal{A}$ is a B -module with a multiplication $\mu : \mathcal{A} \otimes_A \mathcal{A} \rightarrow \mathcal{A}$ which is a homomorphism of A -modules. We say that μ is Δ invariant if

$$b\mu(x \otimes y) = \mu\Delta(b)(x \otimes y) \quad \text{for } b \in B, \quad x, y \in \mathcal{A}, \quad (2.5)$$

and Φ associative, if

$$\mu(\Phi_1 x \otimes \mu(\Phi_2 y \otimes \Phi_3 z)) = \mu(\mu(x \otimes y) \otimes z) \quad \text{for } x, y, z \in \mathcal{A}. \quad (2.6)$$

Note, that a B -module $\mathcal{A}$ equipped with Δ invariant and Φ associative multiplication is an associative algebra in the monoidal category $\mathcal{C}(B, \Delta, \Phi)$. The multiplication $\tilde{\mu} = \mu F^{-1} : M \otimes_A M \rightarrow M$ will be $\tilde{\Phi}$ -associative and invariant in the category $\tilde{\mathcal{C}}$.

We are interested in the case when $A = \mathbb{C}[[h]]$, $B = U(\mathfrak{g})[[h]]$ where $\mathfrak{g}$ is a complex simple Lie algebra. In this case, all tensor products over $\mathbb{C}[[h]]$ are completed in h -adic topology.

Denote by $\varphi \in \wedge^{\otimes 3} \mathfrak{g}$ an invariant element (unique up to scaling for $\mathfrak{g}$ simple) and by $r \in \wedge^{\otimes 2} \mathfrak{g}$ the so-called Drinfeld-Jimbo r -matrix of the form (3.3) such that the Schouten bracket of r with itself is equal to φ :

$$[[r, r]] = \varphi. \quad (2.7)$$

In [Dr1], Drinfeld proved the following (see also [DS2] for the property c)).

Proposition 2.1. *1. There is an invariant element $\Phi_h \in U(\mathfrak{g})[[h]]^{\otimes 3}$ of the form $\Phi_h = 1 \otimes 1 \otimes 1 + h^2 \varphi + \dots$ satisfying the following properties:*

- a) it depends on h^2 , i.e. $\Phi_h = \Phi_{-h}$;*
- b) it satisfies the equations (2.1) (i.e. invariant) and (2.2) with the usual Δ arising from $U(\mathfrak{g})$;*

- c) it is invariant under the Cartan involution θ ;
d) $\Phi_h^{-1} = \Phi_h^{321}$, where $\Phi^{321} = \Phi_3 \otimes \Phi_2 \otimes \Phi_1$ for $\Phi = \Phi_1 \otimes \Phi_2 \otimes \Phi_3$;
2. There is an element $F_h \in U(\mathfrak{g})[[h]]^{\otimes 2}$ of the form $F_h = 1 \otimes 1 + (h/2)r + \dots$ satisfying the equation (2.4) with the usual Δ and with $\tilde{\Phi} = 1 \otimes 1 \otimes 1$.

This proposition implies that there are two nontrivial Drinfeld algebras: the first, $(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ with the usual comultiplication and Φ from Proposition 2.1, and the second, $(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$ where $\tilde{\Delta}(x) = F_h \Delta(x) F_h^{-1}$ for $x \in U(\mathfrak{g})$. The pair (Id, F_h) defines an equivalence between the corresponding monoidal categories $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ and $\mathcal{C}(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$.

It is clear that reduction modulo h defines a functor from either of these categories to the category of representations of $U(\mathfrak{g})$ and the equivalence just described reduces to the identity modulo h . In fact, both categories are $\mathbb{C}[[h]]$ -linear extensions of the $\mathbb{C}$ -linear category of representations of $\mathfrak{g}$. Ignoring the monoidal structure the extension is a trivial one, but the associator Φ in the first case and the comultiplication $\tilde{\Delta}$ in the second case make the extension non-trivial from the point of view of monoidal categories.

The bialgebra $U(\mathfrak{g})[[h]]$ with comultiplication $\Delta_h = \tilde{\Delta}$ is denoted by $U_h(\mathfrak{g})$ and is isomorphic to the Drinfeld-Jimbo quantum group ([Dr1]).

Let $\mathcal{A}$ be a $U(\mathfrak{g})$ invariant commutative algebra, i.e. an algebra with $U(\mathfrak{g})$ invariant multiplication μ in sense of (2.5). A quantization of $\mathcal{A}$ is an associative algebra, $\mathcal{A}_h$, which is isomorphic to $\mathcal{A}[[h]] = \mathcal{A} \otimes \mathbb{C}[[h]]$ (completed tensor product) as a $\mathbb{C}[[h]]$ -module, with multiplication in $\mathcal{A}_h$ having the form $\mu_h = \mu + h\mu_1 + o(h)$. The Poisson bracket corresponding to the quantization is given by $\{a, b\} = \mu_1(a, b) - \mu_1(b, a)$, $a, b \in \mathcal{A}$.

In general, we call a skew-symmetric bilinear form $\mathcal{A} \otimes \mathcal{A} \rightarrow \mathcal{A}$ a bracket, if it satisfies the Leibniz rule in either argument when the other is fixed. The term Poisson bracket indicates that the Jacobi identity is also true. A bracket of the form

$$\{a, b\}_r = (r_1 a)(r_2 b) = \mu(r_1 a, r_2 b) \quad a, b \in \mathcal{A}, \quad (2.8)$$

where $r = r_1 \otimes r_2$ (summation implicit) is the representation of r -matrix r will be called an r -matrix bracket.

Assume, $\mathcal{A}_h$ is a $U_h(\mathfrak{g})$ invariant quantization, i.e. the multiplication μ_h is Δ_h invariant. We shall show that in this case the Poisson bracket $\{\cdot, \cdot\}$ has a special form. Suppose f and g are two brackets on $\mathcal{A}$. Then we define their Schouten bracket $\llbracket f, g \rrbracket$ as

$$\llbracket f, g \rrbracket(a, b, c) = f(g(a, b), c) + g(f(a, b), c) + \text{cyclic permutations of } a, b, c. \quad (2.9)$$

Then $\llbracket f, g \rrbracket$ is a skew-symmetric map $\mathcal{A}^{\otimes 3} \rightarrow \mathcal{A}$. We call f and g compatible, if $\llbracket f, g \rrbracket = 0$.

Proposition 2.2. *Let $\mathcal{A}$ be a $U(\mathfrak{g})$ invariant commutative algebra and $\mathcal{A}_h$ a $U_h(\mathfrak{g})$ invariant quantization. Then the corresponding Poisson bracket has the form*

$$\{a, b\} = f(a, b) - \{a, b\}_r \quad (2.10)$$

where $f(a, b)$ is a $U(\mathfrak{g})$ invariant bracket.

The brackets f and $\{\cdot, \cdot\}_r$ are compatible and $\llbracket f, f \rrbracket = -\varphi_{\mathcal{A}}$ where $\varphi_{\mathcal{A}}(a, b, c) = (\varphi_1 a)(\varphi_2 b)(\varphi_3 c)$ and $\varphi_1 \otimes \varphi_2 \otimes \varphi_3 = \varphi \in \wedge^3 \mathfrak{g}$ is the invariant element.

Proof. The permutation $\sigma : \mathcal{A}_h^{\otimes 2} \rightarrow \mathcal{A}_h^{\otimes 2}$, $a \otimes b \rightarrow b \otimes a$, is an equivariant operator in the category $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$, because Δ is a cocommutative comultiplication. The equivalence of categories $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ and $\mathcal{C}(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$ implies that the operator $\tilde{\sigma} = F\sigma F^{-1}$ on $\mathcal{A}^{\otimes 2}$ is equivariant under action of the quantum group $U_h(\mathfrak{g})$.

Suppose the multiplication in $\mathcal{A}_h$ has the form $\mu_h(a, b) = ab + h\mu_1(a, b) + o(h)$. It is easy to calculate that

$$\frac{1}{h}\mu_h(Id - \tilde{\sigma})(a \otimes b) = \mu_1(a, b) - \mu_1(b, a) + (r_1 a)(r_2 b) + O(h) = \{a, b\} + \{a, b\}_r + O(h).$$

But this is a $U_h(\mathfrak{g})$ equivariant operator $\mathcal{A}_h^{\otimes 2} \rightarrow \mathcal{A}_h$. Taking $h = 0$ we obtain that the bracket $f(a, b) = \{a, b\} + \{a, b\}_r$ must be $U(\mathfrak{g})$ invariant. So, we have $\{a, b\} = f(a, b) - \{a, b\}_r$, as required.

It is easy to check that any bracket of the form $\{a, b\} = (X_1 a)(X_2 b) = \mu(X_1 a, X_2 b)$, for $X_1 \otimes X_2 \in \mathfrak{g} \wedge \mathfrak{g}$, is compatible with any invariant bracket. In particular, an r -matrix bracket is compatible with f . In addition, $\{\cdot, \cdot\}$ is a Poisson bracket, so its Schouten bracket with itself is equal to zero. Using this and taking into account that the Schouten bracket of r -matrix bracket with itself is equal to $\varphi_{\mathcal{A}}$, we obtain from (2.10) that $\llbracket f, f \rrbracket = -\varphi_{\mathcal{A}}$. $\square$

Remark 2.1. a) It is clear that if r satisfies (2.7), then $-r$ satisfies (2.7), too, and we may replace F_h in Proposition 2.1 with $\tilde{F}_h$ with leading terms $1 \otimes 1 - (h/2)r$. Then, instead (2.10) we can write $\{a, b\} = f(a, b) + \{a, b\}_r$.

b) Assume that $\mathcal{A}_{t,h}$ is a two parameter quantization of $\mathcal{A}$, i.e. a topologically free $\mathbb{C}[[t, h]]$ -module with a multiplication of the form $\mu_{t,h}(a, b) = ab + h\mu_1(a, b) + t\mu'_1(a, b) + o(t, h)$. Assume, that $\mathcal{A}_{t,h}$ is $U_h(\mathfrak{g})$ invariant, so that $\mathcal{A}_{t,0}$ is $U(\mathfrak{g})$ invariant. Then there are two compatible Poisson brackets corresponding to such a quantization: the bracket $\mu_1(a, b) - \mu_1(b, a)$ of the form (2.10) and the $U(\mathfrak{g})$ invariant bracket $v(a, b) = \mu'_1(a, b) - \mu'_1(b, a)$. Since v is invariant, the compatibility is equivalent to $\llbracket f, v \rrbracket = 0$.

c) In view of equivalence of the categories $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ and $\mathcal{C}(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$, the problem of quantizing the algebra $\mathcal{A}$ may be considered in the first category. If $\mathcal{A}_h$ is a $U_h(\mathfrak{g})$ invariant quantization with multiplication μ_h , then the multiplication $\bar{\mu}_h = \mu_h F_h = \mu + h\bar{\mu}_1 + o(h)$ will be $U(\mathfrak{g})$ invariant and Φ_h associative in sense of (2.6). We have $\bar{\mu}_1(a, b) - \bar{\mu}_1(b, a) = f(a, b)$, where f is from (2.10). So, we see that the invariant bracket f from (2.10) with $\llbracket f, f \rrbracket = -\varphi_{\mathcal{A}}$ plays the role of Poisson bracket for Φ_h associative quantization. Similarly, the two parameter quantization $\mathcal{A}_{t,h}$ corresponds to the $U(\mathfrak{g})$ invariant Φ_h associative quantization in $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ with a pair of compatible invariant brackets f and v where $\llbracket f, f \rrbracket = -\varphi_{\mathcal{A}}$ and $\llbracket v, v \rrbracket = 0$. Working in the category $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ can simplify the process of quantization. (see [DS1] and Section 5).

In the next section we consider the case when $\mathcal{A}$ is a function algebra on a semisimple orbit in the coadjoint representation of $\mathfrak{g}$ and we give a classification of the invariant brackets f satisfying the property $\llbracket f, f \rrbracket = -\varphi_{\mathcal{A}}$. Moreover, among such f we distinguish those which are compatible with the KKS Poisson brackets.

3 Pairs of brackets on semisimple orbits

Let $\mathfrak{g}$ be a simple complex Lie algebra, $\mathfrak{h}$ a fixed Cartan subalgebra. Let $\Omega \subset \mathfrak{h}^*$ be the system of roots corresponding to $\mathfrak{h}$. Select a system of positive roots, Ω^+ , and denote by $\Pi \subset \Omega$ the subset of simple roots. Fix an element $E_\alpha \in \mathfrak{g}$ of weight α for each $\alpha \in \Omega^+$ and choose $E_{-\alpha}$ such that $(E_\alpha, E_{-\alpha}) = 1$ for the Killing form $(\cdot, \cdot)$ on $\mathfrak{g}$. Then, for all pairs of roots α, β such that $\alpha + \beta \neq 0$ we define the numbers $N_{\alpha, \beta}$ in the following way:

$$\begin{aligned} [E_\alpha, E_\beta] &= N_{\alpha, \beta} E_{\alpha+\beta} & \text{if } \alpha + \beta \in \Omega, \\ N_{\alpha, \beta} &= 0 & \text{if } \alpha + \beta \notin \Omega. \end{aligned}$$

These numbers satisfy the following property, [He]. For the roots α, β, γ such that $\alpha + \beta + \gamma = 0$ one has

$$N_{\alpha, \beta} = N_{\beta, \gamma} = N_{\gamma, \alpha}. \quad (3.1)$$

Let Γ be a subset of Π . Denote by $\mathfrak{h}_\Gamma^*$ the subspace in $\mathfrak{h}^*$ generated by Γ . Note, that $\mathfrak{h}^* = \mathfrak{h}_\Gamma^* \oplus \mathfrak{h}_{\Pi \setminus \Gamma}^*$, and one can identify $\mathfrak{h}_{\Pi \setminus \Gamma}^*$ and $\mathfrak{h}^*/\mathfrak{h}_\Gamma^*$ via the projection $\mathfrak{h}^* \rightarrow \mathfrak{h}^*/\mathfrak{h}_\Gamma^*$.

Let $\Omega_\Gamma \subset \mathfrak{h}_\Gamma^*$ be the subsystem of roots in Ω generated by Γ , i.e. $\Omega_\Gamma = \Omega \cap \mathfrak{h}_\Gamma^*$. Denote by $\mathfrak{g}_\Gamma$ the subalgebra of $\mathfrak{g}$ generated by the elements $\{E_\alpha, E_{-\alpha}\}$, $\alpha \in \Gamma$, and $\mathfrak{h}$. Such a subalgebra is called the Levi subalgebra.

Let G be a complex connected Lie group with Lie algebra $\mathfrak{g}$ and G_Γ a subgroup with Lie algebra $\mathfrak{g}_\Gamma$. Such a subgroup is called the Levi subgroup. It is known that G_Γ is a connected subgroup. Let M be a homogeneous space of G and G_Γ be the stabilizer of a point $o \in M$. We can identify M and the coset space G/G_Γ . It is known, that such M is isomorphic to a semisimple orbit in $\mathfrak{g}^*$. This orbit goes through an element $\lambda \in \mathfrak{g}^*$ which is just the trivial extension to all of $\mathfrak{g}^*$ (identifying $\mathfrak{g}$ and $\mathfrak{g}^*$ via the Killing form) of a map $\lambda : \mathfrak{h}_{\Pi \setminus \Gamma} \rightarrow \mathbb{C}$ such that $\lambda(\alpha) \neq 0$ for all $\alpha \in \Pi \setminus \Gamma$. Conversely, it is easy to show that any semisimple orbit in $\mathfrak{g}^*$ is isomorphic to the quotient of G by a Levi subgroup.

The projection $\pi : G \rightarrow M$ induces the map $\pi_* : \mathfrak{g} \rightarrow T_o$ where T_o is the tangent space to M at the point o . Since the ad-action of $\mathfrak{g}_\Gamma$ on $\mathfrak{g}$ is semisimple, there exists an $\text{ad}(\mathfrak{g}_\Gamma)$ -invariant subspace $\mathfrak{m} = \mathfrak{m}_\Gamma$ of $\mathfrak{g}$ complementary to $\mathfrak{g}_\Gamma$, and one can identify T_o and $\mathfrak{m}$ by means of π_* . It is easy to see that subspace $\mathfrak{m}$ is uniquely defined and has a basis formed by the elements $E_\gamma, E_{-\gamma}$, $\gamma \in \Omega^+ \setminus \Omega_\Gamma$.

Let $v \in \mathfrak{g}^{\otimes m}$ be a tensor over $\mathfrak{g}$. Using the right and the left actions of G on itself, one can associate with v right and left invariant tensor fields on G denoted by v^r and v^l .

We say that a tensor field, t , on G is right G_Γ invariant, if t is invariant under the right action of G_Γ . The G equivariant diffeomorphism between M and G/G_Γ implies that any right G_Γ invariant tensor field t on G induces tensor field $\pi_*(t)$ on M . The field $\pi_*(t)$ will be invariant on M if, in addition, t is left invariant on G , and any invariant tensor field on M can be obtained in such a way. Let $v \in \mathfrak{g}^{\otimes m}$. For v^l to be right G_Γ invariant it is necessary and sufficient that v to be $\text{ad}(\mathfrak{g}_\Gamma)$ invariant. Denote $\pi^r(v) = \pi_*(v^r)$ for any tensor v on $\mathfrak{g}$ and $\pi^l(v) = \pi_*(v^l)$ for any $\text{ad}(\mathfrak{g}_\Gamma)$ invariant tensor v on $\mathfrak{g}$. Note, that tensor $\pi^r(v)$ coincides with the image of v by the map $\mathfrak{g}^{\otimes m} \rightarrow \text{Vect}(M)^{\otimes m}$ induced by the action map $\mathfrak{g} \rightarrow \text{Vect}(M)$. Any G invariant tensor on M has the form $\pi^l(v)$. Moreover, v clearly can be uniquely chosen from $\mathfrak{m}^{\otimes m}$.

Denote by $\llbracket v, w \rrbracket \in \wedge^{k+l-1} \mathfrak{g}$ the Schouten bracket of the polyvectors $v \in \wedge^k \mathfrak{g}$, $w \in \wedge^l \mathfrak{g}$, defined by the formula

$$\llbracket X_1 \wedge \cdots \wedge X_k, Y_1 \wedge \cdots \wedge Y_l \rrbracket = \sum (-1)^{i+j} [X_i, Y_j] \wedge X_1 \wedge \cdots \hat{X}_i \cdots \hat{Y}_j \cdots \wedge Y_l,$$

where $[\cdot, \cdot]$ is the bracket in $\mathfrak{g}$. The Schouten bracket is defined in the same way for polyvector fields on a manifold, but instead of $[\cdot, \cdot]$ one uses the Lie bracket of vector fields. We will use the same notation for the Schouten bracket on manifolds. It is easy to see that $\pi^r(\llbracket v, w \rrbracket) = \llbracket \pi^r(v), \pi^r(w) \rrbracket$, and the same relation is valid for π^l .

Denote by $\overline{\Omega}_\Gamma$ the image of Ω in $\mathfrak{h}_{\Pi \setminus \Gamma}^*$ without zero. It is clear that $\Omega_{\Pi \setminus \Gamma}$ can be identified with a subset of $\overline{\Omega}_\Gamma$ and each element from $\overline{\Omega}_\Gamma$ is a linear combination of elements from $\Pi \setminus \Gamma$ with integer coefficients which are all positive or negative. Thus, the subset $\overline{\Omega}_\Gamma^+ \subset \overline{\Omega}_\Gamma$ of the elements with positive coefficients is exactly the image of Ω^+ . We call elements of $\overline{\Omega}_\Gamma$ quasiroots and the images of $\Pi \setminus \Gamma$ simple quasiroots.

Proposition 3.1. *a) Let β and β' be roots from Ω such that they give the same element in $\overline{\Omega}_\Gamma$. Then there exist roots $\alpha_1, \dots, \alpha_k \in \Omega_\Gamma$ such that $\beta + \alpha_1 + \cdots + \alpha_k = \beta'$ and all partial sums $\beta + \alpha_1 + \cdots + \alpha_i$, $i = 1, \dots, k$ are roots.*

b) Let $\bar{\beta}_1, \dots, \bar{\beta}_k, \bar{\beta}$ be elements of $\overline{\Omega}_\Gamma$ such that $\bar{\beta} = \bar{\beta}_1 + \cdots + \bar{\beta}_k$. Then there exist representatives of these elements in Ω such that $\beta = \beta_1 + \cdots + \beta_k$.

Proof. a) Let $\beta' = \beta + \gamma_1 + \cdots + \gamma_m$ where $\gamma_i \in \Gamma \cup -\Gamma$. If $(\beta', \beta) > 0$, then $\beta' - \beta$ is a root, and the proposition follows. Proceed by induction on m . Since $(\beta', \beta') > 0$, if $(\beta', \beta) \leq 0$, then there exists a γ_i , say γ_m , such that $(\beta', \gamma_m) > 0$, so $\beta' - \gamma_m$ is a root and $\beta' - \gamma_m = \beta + \gamma_1 + \cdots + \gamma_{m-1}$. By induction, the proposition holds for the pair $\beta' - \gamma_m$ and β , i.e. there exist a representation $\beta + \alpha_1 + \cdots + \alpha_{k-1} = \beta' - \gamma_m$ satisfying the proposition. Now, putting $\alpha_k = \gamma_m$, we obtain the required representation of β' .

b) Let $\beta'_1, \dots, \beta'_k, \beta'$ be some representatives of elements $\bar{\beta}_1, \dots, \bar{\beta}_k, \bar{\beta}$ in $\bar{\Omega}$. Then we have the equation $\beta'_1 + \cdots + \beta'_k + \gamma_1 + \cdots + \gamma_m = \beta'$ for some $\gamma_i \in \Gamma \cup -\Gamma$, $i = 1, \dots, m$. If $(\gamma_i, \beta') > 0$, then $\beta' - \gamma_i$ is a root, and we can take $\beta' - \gamma_i$ instead β' . Using iteration, we can regard that all $(\gamma_i, \beta') \leq 0$. Then, there exists a β'_i , say β'_k , such that $(\beta'_k, \beta') > 0$, so $\beta' - \beta'_k$ is a root. Applying induction on k , one can suppose that there are representatives $\beta_1, \dots, \beta_{k-1}$ of $\bar{\beta}_1, \dots, \bar{\beta}_{k-1}$, such that we have an equation $\beta_1 + \cdots + \beta_{k-1} = \beta' - \beta'_k + \gamma_1 + \cdots + \gamma_n$ for some $\gamma_i \in \Gamma \cup -\Gamma$, $i = 1, \dots, n$, and $\tilde{\beta} = \beta_1 + \cdots + \beta_{k-1}$ is a root. If $(\tilde{\beta}, \gamma_i) > 0$, then for some β_j , say β_1 , $(\beta_1, \gamma_i) > 0$, $\beta_1 - \gamma_i$ is a root, and one can replace β_1 by $\beta_1 - \gamma_i$. Repeating this argument we can assume that all $(\tilde{\beta}, \gamma_i) \leq 0$. Then $(\tilde{\beta}, (\beta' - \beta'_k)) \geq 0$, so either $(\tilde{\beta}, \beta') > 0$, and we set $\beta_k = \beta'_k - \sum \gamma_i$, $\beta = \beta'$, or $(\tilde{\beta}, -\beta'_k) > 0$, and we set $\beta_k = \beta'_k$ and $\beta = \beta' + \sum \gamma_i$. In any case, we obtain the required representatives of $\bar{\beta}_1, \dots, \bar{\beta}_k, \bar{\beta}$. $\square$

Remark 3.1. It is obvious that $\mathfrak{m}$ considered as a $\mathfrak{g}_\Gamma$ representation space decomposes into the direct sum of subrepresentations $\mathfrak{m}_{\bar{\beta}}$, $\bar{\beta} \in \overline{\Omega}_\Gamma$, where $\mathfrak{m}_{\bar{\beta}}$ is generated by all the elements E_β , $\beta \in \Omega$, such that the projection of β is equal to $\bar{\beta}$. Part a) of Proposition 3.1 shows that all $\mathfrak{m}_{\bar{\beta}}$ are irreducible. Part b) together with part a) shows that for $\bar{\beta}_1, \bar{\beta}_2 \in \overline{\Omega}_\Gamma$ such that $\bar{\beta}_1 + \bar{\beta}_2 \in \overline{\Omega}_\Gamma$ one has $[\mathfrak{m}_{\bar{\beta}_1}, \mathfrak{m}_{\bar{\beta}_2}] = \mathfrak{m}_{\bar{\beta}_1 + \bar{\beta}_2}$. Using the Killing form, it is easy to see that representations $\mathfrak{m}_{\bar{\beta}}$ and $\mathfrak{m}_{-\bar{\beta}}$ are dual.

Question. Is it true that for $\bar{\beta}_1, \bar{\beta}_2 \in \bar{\Omega}_\Gamma$ such that $\bar{\beta}_1 + \bar{\beta}_2 \in \bar{\Omega}_\Gamma$ the representation $\mathfrak{m}_{\bar{\beta}_1 + \bar{\beta}_2}$ is contained in $\mathfrak{m}_{\bar{\beta}_1} \wedge \mathfrak{m}_{\bar{\beta}_2} \subset \wedge^2 \mathfrak{m}$ with multiplicity one?

Since $\mathfrak{g}_\Gamma$ contains the Cartan subalgebra $\mathfrak{h}$, each $\mathfrak{g}_\Gamma$ invariant tensor over $\mathfrak{m}$ has to be of weight zero. It follows that there are no invariant vectors in $\mathfrak{m}$. Hence, there are no invariant vector fields on M .

Consider the invariant bivector fields on M . From the above, such fields correspond to the $\mathfrak{g}_\Gamma$ invariant bivectors from $\wedge^2 \mathfrak{m}$. Note, that any $\mathfrak{h}$ invariant bivector from $\wedge^2 \mathfrak{m}$ has to be of the form $\sum c(\alpha) E_\alpha \wedge E_{-\alpha}$.

Proposition 3.2. *A bivector $v \in \wedge^2 \mathfrak{m}$ is $\mathfrak{g}_\Gamma$ invariant if and only if it has the form $v = \sum c(\alpha) E_\alpha \wedge E_{-\alpha}$ where the sum runs over $\alpha \in \Omega^+ \setminus \Omega_\Gamma$, and for two roots α, β which give the same element in $\mathfrak{h}^*/\mathfrak{h}_\Gamma^*$ one has $c(\alpha) = c(\beta)$.*

Proof. In view of Proposition 3.1 a), we may assume that $\alpha = \beta + \gamma$ where $\gamma \in \Omega_\Gamma$. Then the coefficient before $E_{\beta+\gamma} \wedge E_{-\beta}$ in $\llbracket E_\gamma, v \rrbracket$ appears from the terms $E_\beta \wedge E_{-\beta}$ and $E_{\beta+\gamma} \wedge E_{-\beta-\gamma}$ in v , and is equal to $N_{\gamma,\beta} c(\beta) + N_{\gamma,-\beta-\gamma} c(\beta + \gamma)$. But from (3.1) follows that $N_{\gamma,\beta} = -N_{\gamma,-\beta-\gamma}$, so if v is invariant under action of E_γ , i.e. $\llbracket E_\gamma, v \rrbracket = 0$, then $c(\beta) = c(\beta + \gamma)$. $\square$

This proposition shows, that coefficients of an invariant element $v = \sum c(\alpha) E_\alpha \wedge E_{-\alpha}$ depend only of the image of α in $\bar{\Omega}_\Gamma^+$, denoted $\bar{\alpha}$, so v can be written in the form $v = \sum c(\bar{\alpha}) E_\alpha \wedge E_{-\alpha}$. Let $v \in \wedge^2 \mathfrak{m}$ be of the form $v = \sum c(\bar{\alpha}) E_\alpha \wedge E_{-\alpha}$ where the sum runs over $\alpha \in \Omega^+ \setminus \Omega_\Gamma$. Denote by θ the Cartan automorphism of $\mathfrak{g}$. Then, v is θ anti-invariant, i.e. $\theta v = -v$. Hence, any $\mathfrak{g}_\Gamma$ invariant bivector is θ anti-invariant. If $v, w \in \wedge^2 \mathfrak{m}$ are $\mathfrak{g}_\Gamma$ invariant, then $\llbracket v, w \rrbracket$ is θ invariant and is of the form $\llbracket v, w \rrbracket = \sum e(\bar{\alpha}, \bar{\beta}) E_{\alpha+\beta} \wedge E_{-\alpha} \wedge E_{-\beta}$ where roots α, β are both negative or both positive and $e(\bar{\alpha}, \bar{\beta}) = -e(-\bar{\alpha}, -\bar{\beta})$. Hence, to calculate $\llbracket v, w \rrbracket$ for such v and w it is sufficient to calculate coefficients $e(\bar{\alpha}, \bar{\beta})$ for positive $\bar{\alpha}$ and $\bar{\beta}$.

Proposition 3.3. *Let $v = \sum c(\alpha) E_\alpha \wedge E_{-\alpha}$, $w = \sum d(\alpha) E_\alpha \wedge E_{-\alpha}$ be elements from $\mathfrak{g} \wedge \mathfrak{g}$. Then for any positive roots $\alpha, \beta, (\alpha + \beta)$ the coefficient by the term $E_{\alpha+\beta} \wedge E_{-\alpha} \wedge E_{-\beta}$ in $\llbracket v, w \rrbracket$ is equal to*

$$\begin{aligned} N_{\alpha,\beta} & (c(\alpha)(d(\beta) - d(\alpha + \beta)) + \\ & c(\beta)(d(\alpha) - d(\alpha + \beta)) - \\ & c(\alpha + \beta)(d(\alpha) + d(\beta))). \end{aligned} \tag{3.2}$$

Proof. Direct computation, see [KRR]. $\square$

Let $r \in \mathfrak{g} \wedge \mathfrak{g}$ be the Drinfeld-Jimbo r -matrix:

$$r = \sum_{\alpha \in \Omega^+} E_\alpha \wedge E_{-\alpha}. \tag{3.3}$$

Then $\llbracket r, r \rrbracket = \varphi$ is an invariant element in $\wedge^3 \mathfrak{g}$. From Proposition 3.2 follows that r reduced modulo $\mathfrak{g} \wedge \mathfrak{g}_\Gamma$ is $\mathfrak{g}_\Gamma$ -invariant. Hence, r and φ define invariant bivector and

three-vector fields on M , $\pi^l(r)$ and $\pi^l(\varphi)$, which we denote by r_M and φ_M . Recall, that we identify invariant tensor fields on M with invariant tensors in $\mathfrak{m}$.

From Propositions 3.3 and 3.1 b) follows that the condition that the Schouten bracket of bivector $v = \sum c(\bar{\alpha})E_\alpha \wedge E_{-\alpha}$ with itself give $K^2\varphi_M$ for a number K is

$$c(\bar{\alpha} + \bar{\beta})(c(\bar{\alpha}) + c(\bar{\beta})) = c(\bar{\alpha})c(\bar{\beta}) + K^2 \quad (3.4)$$

for all the pairs of positive quasiroots $\bar{\alpha}, \bar{\beta}$ such that $\bar{\alpha} + \bar{\beta}$ is a quasiroot.

Given $c(\bar{\alpha})$ and $c(\bar{\beta})$ and assuming that $c(\bar{\alpha}) + c(\bar{\beta}) \neq 0$ we find that

$$c(\bar{\alpha} + \bar{\beta}) = \frac{c(\bar{\alpha})c(\bar{\beta}) + K^2}{c(\bar{\alpha}) + c(\bar{\beta})}. \quad (3.5)$$

Assume, $\bar{\alpha}, \bar{\beta}, \bar{\gamma}$ are positive quasiroots such that $\bar{\alpha} + \bar{\beta}, \bar{\beta} + \bar{\gamma}, \bar{\alpha} + \bar{\beta} + \bar{\gamma}$ are also quasiroots. Then the number $c(\bar{\alpha} + \bar{\beta} + \bar{\gamma})$ can be calculated formally (ignoring possible division by zero) in two ways, using (3.5) for the pair $c(\bar{\alpha}), c(\bar{\beta} + \bar{\gamma})$ on the right hand side and also for the pair $c(\bar{\alpha} + \bar{\beta}), c(\bar{\gamma})$. But it is easy to check that these two ways give the same value of $c(\bar{\alpha} + \bar{\beta} + \bar{\gamma})$. In this sense the system of equations corresponding to (3.5) for all pairs is consistent.

Let us consider this system more carefully. For any positive quasiroot represented as sum of simple quasiroots, $\bar{\alpha} = \sum a_i \bar{\alpha}_i$, define the height $\text{ht}(\bar{\alpha}) = \sum a_i$. In general the coefficient $c(\bar{\alpha})$ for quasiroots of height l can be formally defined by iterating (3.5). Let $\bar{\alpha} = \bar{\alpha}_1 + \dots + \bar{\alpha}_l$ (with possible repetitions) and set $c_i := c(\bar{\alpha}_i)$, then

$$c(\bar{\alpha}) = \frac{c_1 c_2 \dots c_l + K^2 \sum c_{i_1} c_{i_2} \dots c_{i_{l-2}} + \dots}{\sum c_{i_1} c_{i_2} \dots c_{i_{l-1}} + K^2 \sum c_{i_1} c_{i_2} \dots c_{i_{l-3}} + \dots}. \quad (3.6)$$

This expression can be written as

$$K \frac{(\prod_{i=1}^l (c_i + K) + \prod_{i=1}^l (c_i - K))}{(\prod_{i=1}^l (c_i + K) - \prod_{i=1}^l (c_i - K))}.$$

Assumption Let $(\bar{\alpha}_1, \dots, \bar{\alpha}_k)$ be the k -tuple of simple quasiroots. In the following we will assume that the point $(c_1, \dots, c_k) := (c(\bar{\alpha}_1), \dots, c(\bar{\alpha}_k)) \in \mathbb{C}^k$ does not lie on any of the subvarieties defined by the expressions in the denominator of (3.6) and $c_i \neq 0$ for all $i = 1, \dots, k$.

Proposition 3.4. *a) Given a k -tuple of positive numbers $(c_1, \dots, c_k)$ satisfying the Assumption, equation (3.6) uniquely defines numbers $c(\bar{\alpha})$ for all positive quasiroots $\bar{\alpha} = \sum \bar{\alpha}_i$ such that the bivector $v = \sum c(\bar{\alpha})E_\alpha \wedge E_{-\alpha}$ satisfies the condition $\llbracket v, v \rrbracket = K^2\varphi_M$. b) When $K = 0$, the solution described in part a) defines a Poisson bracket on M . If all $c(\bar{\alpha}_i)$ are nonzero (as in Assumption), there exists a linear form $\lambda \in \mathfrak{h}_{\Pi \setminus \Gamma}^*$ such that*

$$c(\bar{\alpha}) = \frac{1}{\lambda(\bar{\alpha})} \quad (3.7)$$

for all quasiroots $\bar{\alpha}$.

Proof. a) Since by assumption the denominator is never zero, equation (3.6) defines consistently $c(\bar{\beta})$ satisfying (3.5) for all positive $\bar{\beta}$.

b) When $K = 0$ (3.4) becomes $c(\bar{\alpha} + \bar{\beta})(c(\bar{\alpha}) + c(\bar{\beta})) = c(\bar{\alpha})c(\bar{\beta})$. Assuming $c(\bar{\alpha}_i) \neq 0$ for the simple quasiroots, equation (3.4) implies $c(\bar{\alpha}) \neq 0$ for all quasiroots, so setting $\lambda(\bar{\alpha}) = 1/c(\bar{\alpha})$, we find that equation (3.4) is equivalent to the equation $\lambda(\bar{\alpha} + \bar{\beta}) = \lambda(\bar{\alpha}) + \lambda(\bar{\beta})$. Thus λ is a linear functional, i.e., an element of $\mathfrak{h}_{\Pi \setminus \Gamma}^*$, which by the construct must be nonzero on all quasiroots. $\square$

Remark 3.2. a) This proposition shows that invariant brackets v on M such that $\llbracket v, v \rrbracket = K^2 \varphi_M$ form a k -dimensional manifold, $\mathcal{X}_K$, which equals $\mathbb{C}^k$ minus the subvarieties defined in the *Assumption*, where k is the number of elements from $\Pi \setminus \Gamma$. Further, it is known that $k = \dim H^2(M)$, [Bo]. If K is regarded as indeterminate, then v forms a $k + 1$ dimensional manifold, $\mathcal{X} \subset \mathbb{C}^k \times \mathbb{C}$, (component $\mathbb{C}$ corresponds to K). Submanifold $\mathcal{X}_0$ corresponds to $K = 0$, i.e. consists of Poisson brackets. It is easy to see that all the Poisson brackets of the type $c(\bar{\alpha}) = 1/\lambda(\bar{\alpha}) \neq 0$ are nondegenerate. Since $\mathcal{X}$ is connected, it follows that almost all brackets v (except an algebraic subset in $\mathcal{X}$ of lesser dimension) are nondegenerate as well.

b) If v defines a Poisson bracket on M , then M is a symplectic manifold and may be realized as an orbit in $\mathfrak{g}^*$ passing through the element λ from (3.7) trivially extended to $\mathfrak{h}^*$, with the KKS bracket.

Now we fix a Poisson bracket $v = \sum (1/\lambda(\bar{\alpha})) E_\alpha \wedge E_{-\alpha}$ where λ is a fixed linear form and describe the invariant brackets $f = \sum c(\bar{\alpha}) E_\alpha \wedge E_{-\alpha}$ which satisfy the conditions

$$\begin{aligned} \llbracket f, f \rrbracket &= K^2 \varphi_M & \text{for } K \neq 0, \\ \llbracket f, v \rrbracket &= 0. \end{aligned} \quad (3.8)$$

An ordered pairs of quasiroots $\bar{\alpha}, \bar{\beta}$ such that $\bar{\alpha} + \bar{\beta}$ is a quasiroot as well will be called an *admissible pair*. Substituting in (3.2) instead $d(\alpha)$ the coefficients of w , we obtain that the condition $\llbracket f, v \rrbracket = 0$ is equivalent to the system of equations for the coefficients of f

$$c(\bar{\alpha})\lambda(\bar{\alpha})^2 + c(\bar{\beta})\lambda(\bar{\beta})^2 = c(\bar{\alpha} + \bar{\beta})\lambda(\bar{\alpha} + \bar{\beta})^2 \quad (3.9)$$

for all admissible pairs $\bar{\alpha}, \bar{\beta}$.

On the other hand, the condition $\llbracket f, f \rrbracket = K^2 \varphi_M$ is equivalent to the system of equations (3.4) for all admissible pairs of quasiroots.

Substituting $c(\alpha + \beta)$ from (3.9) in (3.4) we obtain

$$(c(\bar{\alpha})\lambda(\bar{\alpha})^2 + c(\bar{\beta})\lambda(\bar{\beta})^2)(c(\bar{\alpha}) + c(\bar{\beta})) = c(\bar{\alpha})c(\bar{\beta})(\lambda(\bar{\alpha}) + \lambda(\bar{\beta}))^2 + K^2\lambda(\bar{\alpha} + \bar{\beta})^2.$$

Cancelling terms and extracting the square root, we obtain the equation

$$c(\bar{\beta})\lambda(\bar{\beta}) = c(\bar{\alpha})\lambda(\bar{\alpha}) \pm K\lambda(\bar{\alpha} + \bar{\beta}). \quad (3.10)$$

Substituting $c(\bar{\beta})\lambda(\bar{\beta})$ from (3.10) in (3.9), we obtain

$$c(\bar{\alpha} + \bar{\beta})\lambda(\bar{\alpha} + \bar{\beta}) = c(\bar{\alpha})\lambda(\bar{\alpha}) \pm K\lambda(\bar{\beta}) \quad (3.11)$$

So, the conditions (3.8) on f are equivalent to the system of equations (3.10, 3.11) with the same sign before K for all admissible pairs $\bar{\alpha}$ and $\bar{\beta}$ from $\overline{\Omega}_\Gamma^+$.

We say that an ordered triple of positive quasiroots (not necessarily different) $\bar{\alpha}, \bar{\beta}, \bar{\gamma} \in \overline{\Omega}_\Gamma^+$ is an *admissible triple*, if $\bar{\alpha} + \bar{\beta}$, $\bar{\beta} + \bar{\gamma}$, and $\bar{\alpha} + \bar{\beta} + \bar{\gamma}$ are quasiroots, too.

Lemma 3.1. *Let $\bar{\alpha}, \bar{\beta}, \bar{\gamma}$ be an admissible triple of quasiroots. If $c(\bar{\alpha} + \bar{\beta})\lambda(\bar{\alpha} + \bar{\beta}) = c(\bar{\alpha})\lambda(\bar{\alpha}) + K\lambda(\bar{\beta})$, then $c(\bar{\beta} + \bar{\gamma})\lambda(\bar{\beta} + \bar{\gamma}) = c(\bar{\beta})\lambda(\bar{\beta}) + K\lambda(\bar{\gamma})$, that is, the signs before K in (3.11) for the admissible pairs α, β and β, γ is the same.*

Proof. The admissible pair $\bar{\alpha}, \bar{\beta} + \bar{\gamma}$ gives the equation

$$c(\bar{\alpha})\lambda(\bar{\alpha}) = c(\bar{\beta} + \bar{\gamma})\lambda(\bar{\beta} + \bar{\gamma}) \pm K\lambda(\bar{\alpha} + \bar{\beta} + \bar{\gamma}).$$

The first equation given in the lemma implies that the $+$ sign appears in equation (3.10). These two equations imply

$$c(\bar{\beta})\lambda(\bar{\beta}) - K\lambda(\bar{\alpha} + \bar{\beta}) = c(\bar{\beta} + \bar{\gamma})\lambda(\bar{\beta} + \bar{\gamma}) \pm K\lambda(\bar{\alpha} + \bar{\beta} + \bar{\gamma}).$$

Substituting for $c(\bar{\beta} + \bar{\gamma})\lambda(\bar{\beta} + \bar{\gamma})$ using (3.11), we get

$$c(\bar{\beta})\lambda(\bar{\beta}) - K\lambda(\bar{\alpha} + \bar{\beta}) = c(\bar{\beta})\lambda(\bar{\beta}) \pm K\lambda(\bar{\gamma}) \pm K\lambda(\bar{\alpha} + \bar{\beta} + \bar{\gamma}),$$

where the last two $\pm$ are independent. However if

$$c(\bar{\beta} + \bar{\gamma})\lambda(\bar{\beta} + \bar{\gamma}) = c(\bar{\beta})\lambda(\bar{\beta}) - K\lambda(\bar{\gamma}),$$

we have a contradiction, since either the sign in front of $K\lambda(\bar{\alpha} + \bar{\beta} + \bar{\gamma})$ is positive and $-K\lambda(\bar{\alpha} + \bar{\beta}) = +K\lambda(\bar{\alpha} + \bar{\beta})$, so $0 = \lambda(\bar{\alpha} + \bar{\beta})$, or the sign in front of $K\lambda(\bar{\alpha} + \bar{\beta} + \bar{\gamma})$ is negative, implying $0 = \lambda(\bar{\gamma})$. We conclude that the sign in front of $\lambda(\bar{\gamma})$ must be positive, $\square$

In the situation of the lemma we can express $c(\bar{\gamma})$ in term of $c(\bar{\alpha})$ as

$$c(\bar{\gamma})\lambda(\bar{\gamma}) = c(\bar{\alpha})\lambda(\bar{\alpha}) + K(\lambda(\bar{\alpha} + \bar{\beta}) + \lambda(\bar{\beta} + \bar{\gamma})). \quad (3.12)$$

Now, we consider the pair (M, λ) as an orbit in $\mathfrak{g}^*$ passing through λ with the KKS Poisson bracket

$$v = v_\lambda = \sum_{\alpha \in \Omega^+} \frac{1}{\lambda(\bar{\alpha})} E_\alpha \wedge E_{-\alpha}.$$

Definition 3.1. We call M a *good orbit*, if there exists on M an invariant bracket $f = \sum c(\bar{\alpha})E_\alpha \wedge E_{-\alpha}$ satisfying the conditions (3.8).

Proposition 3.5. *The good semisimple orbits are as follows:*

- a) For $\mathfrak{g}$ of type A_n all semisimple orbits are good.
- b) For all other $\mathfrak{g}$, the orbit M is good if and only if the set $\Pi \setminus \Gamma$ consists of one or two roots which appear in representation of the maximal root with coefficient 1.

Proof. a) In this case the system of quasiroots $\overline{\Omega}_\Gamma$ looks like a system of roots of type A_k for k being equal to the number of elements of $\Pi \setminus \Gamma$. So, the simple quasiroots can be ordered in a sequence $\bar{\beta}_1, \dots, \bar{\beta}_k$ in such a way that all subsequences consisting of three adjacent elements are admissible. Pick an arbitrary value for $c(\bar{\beta}_1)$ and a sign before K in (3.10) for the pair $\bar{\beta}_1$ and $\bar{\beta}_2$. Then, due to Lemma 3.1, consistency of system (3.10, 3.11) implies that the sign before K is the same for all adjacent pairs $\bar{\beta}_i$ and $\bar{\beta}_{i+1}$. Using equations (3.10) and (3.11) for a fixed sign before K and induction on $\text{ht}(\alpha)$, we find all the coefficients $c(\bar{\alpha})$ of f and see that the system (3.10, 3.11) is consistent.

b) Let $\mathfrak{g}$ be of type B_n or C_n . Then the maximal root has the form $\alpha_1 + 2\alpha_2 + \dots + 2\alpha_n$ where $\alpha_i \in \Pi$. Denote $\beta = \alpha_2 + \dots + \alpha_n$ which is a root. If $\Pi \setminus \Gamma$ does not contain α_1 , i.e. $\bar{\alpha}_1 = 0$, then $\overline{\Omega}_\Gamma^+$ contains the admissible pair $\bar{\beta}, \bar{\beta}$. So, from (3.10) follows that $c(\bar{\beta})\lambda(\bar{\beta}) = c(\bar{\beta})\lambda(\bar{\beta}) \pm 2K\lambda(\bar{\beta})$, i.e. $\lambda(\bar{\beta}) = 0$ which is impossible.

Assume that $\Pi \setminus \Gamma$ contains α_1 and some roots α_i for $i > 1$. Then both $\bar{\alpha}_1$ and $\bar{\beta}$ are not equal to zero, and $\overline{\Omega}_\Gamma^+$ contains the admissible triple $\bar{\beta}, \bar{\alpha}_1, \bar{\beta}$. It follows from (3.12) that $c(\bar{\beta})\lambda(\bar{\beta}) = c(\bar{\beta})\lambda(\bar{\beta}) + 2K\lambda(\bar{\beta} + \bar{\alpha}_1)$, i.e. $\lambda(\bar{\beta} + \bar{\alpha}_1) = 0$ which is impossible as well, because $\bar{\beta} + \bar{\alpha}_1$ is a quasiroot.

So, for consistency of system (3.10, 3.11) in cases B_n and C_n , the set Γ has to contain all the roots α_i , $i > 1$. But in the latter case the system is trivially consistent, because in that case the set of quasiroots looks like A_1 . The homogeneous space G/G_Γ is a symmetric space.

Consider the case D_n .

The maximal root has the form $\alpha_1 + \alpha_2 + \alpha_3 + 2\alpha_4 + \dots + 2\alpha_n$. Denote $\beta = \alpha_4 + \dots + \alpha_n$ which is a root.

Consider several cases.

The cases when two of $\bar{\alpha}_i$, $i \leq 3$, are equal to zero and $\bar{\beta}$ is not equal to zero lead to an inconsistency in the system (3.10, 3.11) in the same way as in the cases B_n and C_n considered above.

Assume, that two of $\bar{\alpha}_i$, $i \leq 3$, say $\bar{\alpha}_1, \bar{\alpha}_3$, and $\bar{\beta}$ are not equal to zero. Then the sequence

$$\bar{\alpha}_1, \bar{\alpha}_2 + \bar{\beta}, \bar{\alpha}_3, \bar{\alpha}_1 + \bar{\beta} \quad (3.13)$$

is a sequence of four nonzero quasiroots. It is easy to see that the subsequences $\bar{\alpha}_1, \bar{\alpha}_2 + \bar{\beta}, \bar{\alpha}_3$ and $\bar{\alpha}_2 + \bar{\beta}, \bar{\alpha}_3, \bar{\alpha}_1 + \bar{\beta}$ form admissible triples in $\overline{\Omega}_\Gamma^+$. From Lemma 3.1 it follows that the sign before K must be the same in (3.10) for all adjacent pairs. Taking, for example, the sign plus and applying (3.12) to the second triple, we obtain the equation

$$c(\bar{\alpha}_1 + \bar{\beta})\lambda(\bar{\alpha}_1 + \bar{\beta}) = c(\bar{\alpha}_2 + \bar{\beta})\lambda(\bar{\alpha}_2 + \bar{\beta}) + K(\lambda(\bar{\alpha}_2 + \bar{\beta} + \bar{\alpha}_3) + \lambda(\bar{\alpha}_3 + \bar{\alpha}_1 + \bar{\beta})). \quad (3.14)$$

From (3.10) applied to the pair $\bar{\alpha}_1, \bar{\alpha}_2 + \bar{\beta}$ we obtain

$$c(\bar{\alpha}_2 + \bar{\beta})\lambda(\bar{\alpha}_2 + \bar{\beta}) = c(\bar{\alpha}_1)\lambda(\bar{\alpha}_1) + K\lambda(\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\beta}). \quad (3.15)$$

Putting $c(\bar{\alpha}_1)$ from (3.15) in (3.14) and taking into account linearity of λ , we obtain the equality

$$c(\bar{\alpha}_1 + \bar{\beta})\lambda(\bar{\alpha}_1 + \bar{\beta}) = c(\bar{\alpha}_1)\lambda(\bar{\alpha}_1) + 2K\lambda(\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\alpha}_3 + \bar{\beta}) + K\lambda(\bar{\beta}). \quad (3.16)$$

On the other hand, expressing $c(\bar{\alpha}_1 + \bar{\beta})$ in terms of $c(\bar{\alpha}_1)$ from (3.11), we obtain

$$c(\bar{\alpha}_1 + \bar{\beta})\lambda(\bar{\alpha}_1 + \bar{\beta}) = c(\bar{\alpha}_1)\lambda(\bar{\alpha}_1) \pm K\lambda(\bar{\beta}). \quad (3.17)$$

Now, comparing (3.16) and (3.17), we see that if we take plus before K in (3.17), then $\lambda(\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\alpha}_3 + 2\bar{\beta}) = 0$, if we take minus, then $\lambda(\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\alpha}_3 + \bar{\beta}) = 0$. But both of the cases are impossible, since λ is not equal to zero on quasiroots.

Next, assume that $\bar{\beta} = 0$ but $\bar{\alpha}_i \neq 0$ for $i = 1, 2, 3$. In this case the sequence (3.13) makes into the sequence $\bar{\alpha}_1, \bar{\alpha}_2, \bar{\alpha}_3, \bar{\alpha}_1$. Using the above arguments, we obtain that in this case must be $\lambda(\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\alpha}_3) = 0$ which is impossible, since $\bar{\alpha}_1 + \bar{\alpha}_2 + \bar{\alpha}_3$ is a quasiroot.

In the case when $\Pi \setminus \Gamma$ contains only one or two roots of α_i , $i = 1, 2, 3$, system (3.10, 3.11) is consistent, because in these cases the set of quasiroots $\bar{\Omega}_\Gamma$ looks like the system of roots of type A_1 or A_2 .

So, the proposition is proved for the classical $\mathfrak{g}$. For the exceptional $\mathfrak{g}$ the proposition follows from the same arguments. $\square$

Remark 3.3. a) Note that Proposition 3.5 may be reformulated in the following way:

An orbit M is good if and only if the corresponding system of quasiroots $\bar{\Omega}_\Gamma$ is isomorphic to a system of roots of type A_k . We say in this case that M is of type A_k .

Orbits of type A_1 are exactly the orbits which are symmetric spaces. For such orbits $\varphi_M = 0$, and we may take $f = 0$. Symmetric orbits exist for all classical $\mathfrak{g}$ and also for $\mathfrak{g}$ of types E_6, E_7 .

Orbits of type A_2 exist for $\mathfrak{g}$ of type A_n, D_n , and E_6 .

Orbits of type A_k , $k > 2$, exist only in case $\mathfrak{g} = sl(n)$. Moreover, in this case all semisimple orbits have the type A_k for $k \geq 1$.

b) From the proof of Proposition 3.5 it follows that the bracket f satisfying (3.8) is defined on good orbits by the value of its coefficient $c(\alpha)$ for a fixed simple root and the choice of a sign before K . On the other hand, if a fixed f_0 satisfies (3.8), then the family $\pm f_0 + sv$ for arbitrary numbers s also satisfies these conditions. So, this family consists of all invariant brackets satisfying (3.8). Almost all brackets from this family (except a finite number) are nondegenerate, since v is nondegenerate and, therefore, for large s_0 the bracket $\pm f_0 + s_0 v$ is nondegenerate as well.

For symmetric orbits $\Pi \setminus \Gamma$ consists of one element, there is one quasiroot and f_0 is a multiple of v . Note, that in [GP] is given a classification of all orbits in coadjoint representation (not necessarily semisimple) for which $\varphi_M = 0$.

In particular, if we take $K = 1$ and find f_0 such that $\llbracket f_0, f_0 \rrbracket = \varphi_M$, then the family $\pm i f_0 + sw$ gives all the brackets satisfying $\llbracket f_0, f_0 \rrbracket = -\varphi_M$ and compatible with KKS bracket on M .

Note, that if f is a bracket satisfying $\llbracket f, f \rrbracket = -\varphi_M$ and $\{\cdot, \cdot\}_r$ is the r -matrix bracket (2.8), then $f \pm \{\cdot, \cdot\}_r$ is a Poisson bracket on M compatible with KKS bracket.

4 Cohomologies defined by invariant brackets

In the next section we prove the existence of a $U_h(\mathfrak{g})$ invariant quantization of the Poisson brackets described above using the methods of [DS1]. This requires us to consider the

3-cohomology of the complex $(\Lambda^\bullet(\mathfrak{g}/\mathfrak{g}_\Gamma))^{\mathfrak{g}_\Gamma} = (\Lambda^\bullet \mathfrak{m})^{\mathfrak{g}_\Gamma}$ of $\mathfrak{g}_\Gamma$ invariants with differential given by the Schouten bracket with the bivector $v \in (\Lambda^2 \mathfrak{m})^{\mathfrak{g}_\Gamma}$ from Proposition 3.4 a),

$$\delta_v : u \mapsto \llbracket v, u \rrbracket \quad \text{for } u \in (\Lambda^\bullet \mathfrak{m})^{\mathfrak{g}_\Gamma}.$$

The condition $\delta_v^2 = 0$ follows from the Jacobi identity for the Schouten bracket together with the fact that $\llbracket v, v \rrbracket = K^2 \varphi_M$. The latter equation is equivalent to $\llbracket v, v \rrbracket = \varphi$ modulo $\mathfrak{g}_\Gamma \wedge \mathfrak{g} \wedge \mathfrak{g}$, hence $\llbracket v, v \rrbracket$ is invariant modulo $\mathfrak{g}_\Gamma \wedge \mathfrak{g} \wedge \mathfrak{g}$. Denote these cohomologies by $H^k(M, \delta_v)$, whereas the usual de Rham cohomologies are denoted by $H^k(M)$.

Recall, Remark 3.2 a), that the brackets v satisfying $\llbracket v, v \rrbracket = K^2 \varphi$ form a connected manifold $\mathcal{X}$ which contains a submanifold $\mathcal{X}_0$ of Poisson brackets.

Proposition 4.1. *For almost all $v \in \mathcal{X}$ (except an algebraic subset of lesser dimension) one has*

$$H^k(M, \delta_v) = H^k(M)$$

for all k . In particular, $H^k(M, \delta_v) = 0$ for odd k .

Proof. First, let v be a Poisson bracket, i.e. $v \in \mathcal{X}_0$. Then the complex of polyvector fields on M , $\Theta^\bullet$, with the differential δ_v is well defined. Denote by $\Omega^\bullet$ the de Rham complex on M . Since none of the coefficients $c(\bar{\alpha})$ of v are zero, v is a nondegenerate bivector field, and therefore it defines an $\mathcal{A}$ -linear isomorphism $\tilde{v} : \Omega^1 \rightarrow \Theta^1$, $\omega \mapsto v(\omega, \cdot)$, which can be extended up to the isomorphism $\tilde{v} : \Omega^k \rightarrow \Theta^k$ of k -forms onto k -vector fields for all k . Using Jacobi identity for v and invariance of v , one can show that $\tilde{v}$ gives a G invariant isomorphism of these complexes, so their cohomologies are the same.

Since $\mathfrak{g}$ is simple, the subcomplex of $\mathfrak{g}$ invariants, $(\Omega^\bullet)^\mathfrak{g}$, splits off as a subcomplex of $\Omega^\bullet$. In addition, $\mathfrak{g}$ acts trivially on cohomologies, since for any $g \in G$ the map $X \rightarrow X$, $x \mapsto gx$, is homotopic to the identity map. (G is assumed connected.) It follows that cohomologies of complexes $(\Omega^\bullet)^\mathfrak{g}$ and $\Omega^\bullet$ coincide.

But $\tilde{v}$ gives an isomorphism of complexes $(\Omega^\bullet)^\mathfrak{g}$ and $(\Theta^\bullet)^\mathfrak{g} = ((\Lambda^\bullet \mathfrak{m})^{\mathfrak{g}_\Gamma}, \delta_v)$. So, cohomologies of the latter complex coincide with de Rham cohomologies, which proves the proposition for v being Poisson brackets.

Now, consider the family of complexes $((\Lambda^\bullet \mathfrak{m})^{\mathfrak{g}_\Gamma}, \delta_v)$, $v \in \mathcal{X}$. It is clear that δ_v depends algebraically on v . It follows from the uppersemicontinuity of $\dim H^k(M, \delta_v)$ and the fact that $H^k(M) = 0$ for odd k , [Bo], that $H^k(M, \delta_v) = 0$ for odd k and almost all $v \in \mathcal{X}$. Using the uppersemicontinuity again and the fact that the number $\sum_k (-1)^k \dim H^k(M, \delta_v)$ is the same for all $v \in \mathcal{X}$, we conclude that $\dim H^k(M, \delta_v) = \dim H^k(M)$ for even k and almost all v . $\square$

Remark 4.1. Call $v \in \mathcal{X}$ admissible, if it satisfies Proposition 4.1. From the proof of the proposition follows that the subset $\mathcal{D}$ such that $\mathcal{X} \setminus \mathcal{D}$ consists of admissible brackets does not intersect with the subset $\mathcal{X}_0$ consisting of Poisson brackets.

Let M be a good orbit and $f_0 + sw$ the family from Remark 3.3 b) satisfying (3.8) for a fixed K . Then for almost all numbers s this bracket is admissible. Indeed, this family is contained in the two parameter family $tf_0 + sw$. By $t = 0$, $s \neq 0$ we obtain admissible brackets. So, there exist $t_0 \neq 0$ and s_0 such that the bracket $t_0 f_0 + s_0 w$ is admissible. It

follows that the bracket $f_0 + (s_0/t_0)w$ is admissible, too. So, in the family $f_0 + sw$ there is an admissible bracket, and we conclude that almost all brackets in this family (except a finite number) are admissible.

Question. Is it true that the set of admissible brackets contains all the nondegenerate brackets?

For the proof of existence of two parameter quantization for the cases D_n and E_6 in the next section we will use the following result on invariant three-vector fields.

Denote by θ the Cartan automorphism of $\mathfrak{g}$.

Lemma 4.1. *For either D_n or E_6 and one of the subsets, Γ , of simple roots such that G_Γ defines a good orbit, any $\mathfrak{g}_\Gamma$ and θ invariant element v in $\Lambda^3 \mathfrak{m}$ is a multiple of φ_M , that is,*

$$(\Lambda^3(\mathfrak{m})^{\mathfrak{g}_\Gamma} \cong \langle \varphi_M \rangle.$$

Proof. Let $\mathfrak{g}$ be a simple Lie algebra of type D_n or E_6 and $\{\alpha_1, \dots, \alpha_n\}$ a system of simple roots. Changing notation slightly from Section 3, we assume that for $\mathfrak{g} = D_n$, $(\alpha_i, \alpha_{i+1}) = -1$ for $i = 1, \dots, n-2$, $(\alpha_{n-2}, \alpha_n) = -1$ with all other inner products of distinct simple roots are zero, and for $\mathfrak{g} = E_6$, the non-zero products are $(\alpha_i, \alpha_{i+1}) = -1$ for $i = 1, 2, 3, 4$ and $(\alpha_3, \alpha_6) = -1$. For $\mathfrak{g} = D_n$, Γ is one of the subsets of simple roots, $\Gamma_1 = \{\alpha_1, \dots, \alpha_{n-2}\}$, $\Gamma_2 = \{\alpha_2, \dots, \alpha_{n-1}\}$, or $\Gamma_3 = \{\alpha_2, \dots, \alpha_{n-2}, \alpha_n\}$. For $\mathfrak{g} = E_6$, $\Gamma = \{\alpha_2, \alpha_3, \alpha_4, \alpha_6\}$.

The positive quasiroots consist of three elements, $\bar{\alpha}$, $\bar{\alpha}'$, and $\bar{\alpha} + \bar{\alpha}'$. Since a θ invariant element has the form $w + \theta w$ for $w \in \mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'} \otimes \mathfrak{m}_{-(\bar{\alpha} + \bar{\alpha}')}$, It is sufficient to show that the space of invariants in $\mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'} \otimes \mathfrak{m}_{-(\bar{\alpha} + \bar{\alpha}')}$ has dimension one. We know from Remark 3.1 that the subspaces $\mathfrak{m}_{\bar{\alpha}}$, $\mathfrak{m}_{\bar{\alpha}'}$, $\mathfrak{m}_{\bar{\alpha} + \bar{\alpha}'}$ are irreducible representations of $\mathfrak{g}_\Gamma$ and that $\mathfrak{m}_{\bar{\alpha}}$ and $\mathfrak{m}_{-\bar{\alpha}}$ are dual. Therefore the dimension of the space of invariants in $\mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'} \otimes \mathfrak{m}_{-(\bar{\alpha} + \bar{\alpha}')}$ is the multiplicity of the representation $\mathfrak{m}_{\bar{\alpha} + \bar{\alpha}'}$ in the tensor product $\mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'}$. For D_n and any of Γ_i the algebra $\mathfrak{g}_\Gamma \cong A_{n-2}$. For Γ_1 , $\alpha = \alpha_{n-1}$ and $\alpha' = \alpha_n$, the representations $\mathfrak{m}_{\bar{\alpha}}$ and $\mathfrak{m}_{\bar{\alpha}'}$ are both isomorphic to the dual vector representation for A_{n-2} , that is, the contragredient representation to the representation for the fundamental weight λ_{n-2} ,

$$\mathfrak{m}_{\bar{\alpha}_{n-1}} \cong \mathfrak{m}_{\bar{\alpha}_n} \cong (V^{\lambda_{n-2}})^* \cong V^{\lambda_1}.$$

To see that this is so, note first of all that $\mathfrak{m}_{\bar{\alpha}_{n-1}}$ is a lowest weight representation because it has a cyclic vector $E_{\alpha_{n-1}}$ and all negative simple root vectors of $\mathfrak{g}_\Gamma$ annihilate $E_{\alpha_{n-1}}$. The corresponding weight of A_{n-2} is $-\lambda_{n-2}$ because $(\alpha_{n-1}, \alpha_j) = -(\lambda_{n-2}, \alpha_j)$ if $1 \leq j \leq n-2$. The irreducible lowest weight representations with lowest weight $-\lambda_{n-2}$ is $(V^{\lambda_{n-2}})^*$ and A_{n-2} , $(V^{\lambda_j})^* \cong V^{\lambda_{n-1-j}}$. Since the subspaces $\mathfrak{m}_{\bar{\alpha}}$ and $\mathfrak{m}_{\bar{\alpha}'}$ of $\mathfrak{g}$ have zero intersection the wedge product $\mathfrak{m}_{\bar{\alpha}} \wedge \mathfrak{m}_{\bar{\alpha}'}$ projects isomorphically onto the tensor product which contains the representation

$$\mathfrak{m}_{\bar{\alpha} + \bar{\alpha}'} \cong (V^{2\lambda_{n-2}})^* \cong V^{2\lambda_1}$$

with multiplicity one.

In the cases Γ_2 or Γ_3 the representation $\mathfrak{m}_{\bar{\alpha}_1}$ is the contragredient representation to the vector representation $(V^{\lambda_1})^* \cong V^{\lambda_{n-2}}$. The representations $\mathfrak{m}_{\alpha'} = \mathfrak{m}_{\alpha_n}$ for the case Γ_2 and $\mathfrak{m}_{\alpha'} = \mathfrak{m}_{\alpha_{n-1}}$ for Γ_3 are $(V^{\lambda_{n-3}})^* \cong V^{\lambda_2}$. From the elementary representation theory

of the Lie algebra $sl(n-1)$ (type A_{n-2}) we see that tensor product $\mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'}$ contains $\mathfrak{m}_{\bar{\alpha}+\bar{\alpha}'}$ with multiplicity one.

In the case of E_6 , $\mathfrak{g}_\Gamma \cong D_4 \cong so(8)$. The representations $\mathfrak{m}_{\bar{\alpha}}$, $\mathfrak{m}_{\bar{\alpha}'}$ and $\mathfrak{m}_{-(\bar{\alpha}+\bar{\alpha})'}$ are the three inequivalent irreducible eight dimensional representations, the two spinor representations and the vector representation. It is well known that there is a one dimensional space of invariants in the tensor product $\mathfrak{m}_{\bar{\alpha}} \otimes \mathfrak{m}_{\bar{\alpha}'} \otimes \mathfrak{m}_{\bar{\alpha}+\bar{\alpha}'}$. $\square$

The proof of the lemma gives a positive answer on the *Question* from Remark 3.1 for the particular case.

Note that in case of symmetric orbits the space of invariant three-vector fields is equal to zero.

5 $U_h(\mathfrak{g})$ invariant quantizations in one and two parameters

In this section we prove the existence of two types of $U_h(\mathfrak{g})$ invariant quantization of the function algebra $\mathcal{A}$ on $M = G/G_\Gamma$. The first is a one parameter quantization

$$\mu_h(a, b) = \sum_{n \geq 0} h^n \mu_n(a, b) = ab + \sum_{n \geq 1} h^n \mu_n(a, b), \quad \mu_1(a, b) = \frac{1}{2}(f(a, b) - \{a, b\}_r),$$

as described in Section 2, where f is one of the invariant brackets found above,

$$f \in (\Lambda^2 \mathfrak{m})^{\mathfrak{g}}, \quad \llbracket f, f \rrbracket = -\varphi_M.$$

The second is a two parameter quantization

$$\mu_{t,h}(a, b) = ab + h\mu_1(a, b) + t\mu'_1(a, b) + \sum_{k,l \geq 1} h^k t^l \mu_{k,l}(a, b).$$

Recall that in this case there are two compatible Poisson brackets corresponding to such a quantization: the bracket $\mu_1(a, b) - \mu_1(b, a)$ is skew-symmetric of the form (2.10) and $\mu'_1(a, b) - \mu'_1(b, a)$ is a $U(\mathfrak{g})$ invariant bracket $v(a, b)$, which we will assume to be the KKS bracket defined by identifying G/G_Γ with an orbit of the coadjoint representation.

We remind the reader of the method in [DS1]. The first step is to construct a $U(\mathfrak{g})$ invariant quantization in the category $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$. Then we use the equivalence given by the pair (Id, F_h) between the monoidal categories $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ and $\mathcal{C}(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$ to define a $U_h(\mathfrak{g})$ invariant quantization, either $\mu_h F_h^{-1}$ in the one parameter case or $\mu_{t,h} F_h^{-1}$ in the two parameter case (see Section 2). In the first step we used the fact that $(\Lambda^3 \mathfrak{m})^{\mathfrak{g}_\Gamma} = 0$ for symmetric spaces. In the examples considered in this paper, $(\Lambda^3 \mathfrak{m})^{\mathfrak{g}_\Gamma}$ does not necessarily vanish, and we modify the proof using a method from [DS2] (see also [NV]). In the case of A_n any semisimple orbit is a good orbit and a different method is required (see [Do] where the existence of two parameter quantization for maximal orbits is proven). For the cases B_n and C_n , the only good orbits are symmetric spaces and the quantization was dealt with in [DS1]. In the remaining cases $\mathfrak{g} = D_n$ or E_6 , we proved in Lemma 4.1 that $(\Lambda^3 \mathfrak{m})^{\mathfrak{g}_\Gamma, \theta} = \langle \varphi_{\mathcal{A}} \rangle$, and a suitable modification of the proof still applies.

Proposition 5.1. *For almost all (in sense of Proposition 4.1) $\mathfrak{g}$ invariant bracket satisfying $\llbracket f, f \rrbracket = -\varphi_M$, there exists a multiplication μ_h on $\mathcal{A}$*

$$\mu_h(a, b) = ab + (h/2)f(a, b) + \sum_{n \geq 2} h^n \mu_n(a, b)$$

which is $\mathfrak{g}$ invariant (equation 2.5)) and Φ associative (equation (2.6)).

Proof. To begin, consider the multiplication $\mu^{(1)}(a, b) = ab + (h/2)f(a, b)$. The corresponding obstruction cocycle is given by

$$obs_2 = \frac{1}{h^2}(\mu^{(1)}(\mu^{(1)} \otimes id) - \mu^{(1)}(id \otimes \mu^{(1)})\Phi)$$

considered modulo terms of order h . No $\frac{1}{h}$ terms appear because f is a biderivation and, therefore, a Hochschild cocycle. The fact that the presence of Φ does not interfere with the cocycle condition and that this equation defines a Hochschild 3-cocycle was proven in [DS1]. It is well known that if we restrict to the subcomplex of cochains given by differential operators, the differential Hochschild cohomology of $\mathcal{A}$ in dimension p is the space of polyvector fields on M . Since $\mathfrak{g}$ is reductive, the subspace of $\mathfrak{g}$ invariants splits off as a subcomplex and has cohomology given by $(\Lambda^p \mathfrak{m})^{\mathfrak{gr}}$. The complete antisymmetrization of a p -tensor projects the space of invariant differential p -cocycles onto the subspace $(\Lambda^p \mathfrak{m})^{\mathfrak{gr}}$ representing the cohomology. The equation $\llbracket f, f \rrbracket + \varphi_M = 0$ implies that obstruction cocycle is a coboundary, and we can find a 2-cochain μ_2 so that $\mu^{(2)} = \mu^{(1)} + h^2 \mu_2$ satisfies

$$\mu^{(2)}(\mu^{(2)} \otimes id) - \mu^{(2)}(id \otimes \mu^{(2)})\Phi = 0 \text{ mod } h^2.$$

Assume we have defined the deformation $\mu^{(n)}$ to order h^n such that Φ associativity holds modulo h^n , then we define the $(n+1)^{\text{st}}$ obstruction cocycle by

$$obs_{n+1} = \frac{1}{h^{n+1}}(\mu^{(n)}(\mu^{(n)} \otimes id) - \mu^{(n)}(id \otimes \mu^{(n)})\Phi) \text{ mod } h.$$

In [DS1] (Proposition 4) we showed that the usual proof that the obstruction cochain satisfies the cocycle condition carries through to the Φ associative case. The coboundary of obs_{n+1} appears as the h^{n+1} coefficient of the signed sum of the compositions of $\mu^{(n+1)}$ with obs_{n+1} . The fact that $\Phi = 1 \text{ mod } h^2$ together with the pentagon identity implies that the sum vanishes identically, and thus all coefficients vanish, including the coboundary in question. Let $obs'_{n+1} \in (\Lambda^3 \mathfrak{m})^{\mathfrak{gr}}$ be the projection of obs_{n+1} on the totally skew symmetric part, which represents the cohomology class of the obstruction cocycle. The coefficient of h^{n+2} in the same signed sum, when projected on the skew symmetric part is $\llbracket f, obs'_{n+1} \rrbracket$ which is the coboundary of obs'_{n+1} in the complex $(\Lambda^\bullet \mathfrak{m})^{\mathfrak{gr}}, \delta_f = \llbracket f, \cdot \rrbracket)$. Thus obs'_{n+1} is a δ_f cocycle. We have shown in Section 4 that this complex has zero cohomology. Now we modify $\mu^{(n+1)}$ by adding a term $h^n \mu_n$ with $\mu_n \in (\Lambda^2 \mathfrak{m})^{\mathfrak{gr}}$ and consider the $(n+1)^{\text{st}}$ obstruction cocycle for $\mu'^{(n+1)} = \mu^{(n+1)} + h^n \mu_n$. Since the term we added at degree h^n is a Hochschild cocycle we do not introduce a h^n term in the calculation of $\mu^{(n)}(\mu^{(n)} \otimes id) - \mu^{(n)}(id \otimes \mu^{(n)})\Phi$ and the totally skew symmetric projection h^{n+1} term has been modified by $\llbracket f, \mu_n \rrbracket$. By choosing μ_n appropriately we can make the $(n+1)^{\text{st}}$

obstruction cocycle represent the zero cohomology class, and we are able to continue the recursive construction of the desired deformation. $\square$

Now we prove the existence of a two parameter deformation for good orbits in the cases D_n and E_6 .

Proposition 5.2. *Given a pair of $\mathfrak{g}$ invariant brackets, f, v , on a good orbit in D_n or E_6 satisfying $\llbracket f, f \rrbracket = -\varphi_M$, $\llbracket f, v \rrbracket = \llbracket v, v \rrbracket = 0$, there exists a multiplication $\mu_{h,t}$ on $\mathcal{A}$*

$$\mu_{t,h}(a, b) = ab + (h/2)f(a, b) + (t/2)v(a, b) + \sum_{k,l \geq 1} h^k t^l \mu_{k,l}(a, b)$$

which is $\mathfrak{g}$ invariant (equation 2.5)) and Φ associative (equation (2.6)).

Proof. The existence of a multiplication which is Φ associative up to and including h^2 terms is nearly identical to the previous proof. Both f and v are anti-invariant under the Cartan involution θ . We shall look for a multiplication $\mu_{t,h}$ such that $\mu_{k,l}$ is θ anti-invariant and skew-symmetric for odd $k + l$ and θ invariant and symmetric for even $k + l$.

So suppose we have a multiplication defined to order n ,

$$\mu_{t,h}(a, b) = ab + h\mu_1(a, b) + t\mu'_1(a, b) + \sum_{k+l \leq n} h^k t^l \mu_{k,l}(a, b),$$

with mentioned above invariance properties and Φ associative to order h^n .

Using properties c) and d) for Φ from Proposition 2.1, direct computation shows that the obstruction cochain,

$$obs_{n+1} = \sum_{k=0, \dots, n+1} h^k t^{n+1-k} \beta_k,$$

has the following invariance properties: For odd n , obs_{n+1} is θ invariant and $obs_{n+1}(a, b, c) = -obs_{n+1}(c, b, a)$, and for even n , and obs_{n+1} is θ anti-invariant and $obs_{n+1}(a, b, c) = obs_{n+1}(c, b, a)$.

Hence, the projection of obs_{n+1} on $(\Lambda^3 \mathfrak{m})^{\mathfrak{gr}}$ is equal to zero for even n . It follows that all the β_k are Hochschild coboundaries, and the standard argument implies that the multiplication can be extended up to order $n + 1$ with the required properties.

For odd n , Lemma 4.1 shows that the projection on $(\Lambda^3 \mathfrak{m})^{\mathfrak{gr}}$ has the form

$$obs_{n+1} = \left(\sum_{k=0, \dots, n+1} a_k h^k t^{n+1-k} \right) \varphi_M.$$

The KKS bracket is given by the two-vector

$$v = \sum_{\alpha \in \Omega^+ \setminus \Omega_\Gamma} \frac{1}{\lambda(\bar{\alpha})} E_\alpha \wedge E_{-\alpha}.$$

Setting

$$w = \sum_{\alpha \in \Omega^+ \setminus \Omega_\Gamma} \lambda(\bar{\alpha}) E_\alpha \wedge E_{-\alpha},$$

gives

$$\llbracket v, w \rrbracket = -3\varphi_M.$$

Defining

$$\mu'^{(n)} = \mu^{(n)} + \frac{a_0}{3} t^n w,$$

the new obstruction cohomology class is

$$obs'_{n+1} = \left(\sum_{k=1, \dots, n+1} a_k h^k t^{n+1-k} \right) \varphi_M.$$

Finally we define

$$\mu''^{(n)} = \mu'^{(n)} + \sum_{k=1, \dots, n} a_k h^{k-1} t^{n+1-k} f$$

and get an obstruction cocycle which is zero in cohomology. Now the standard argument implies that the deformation can be extended to give a Φ associative invariant multiplication with the required properties of order $n + 1$.

So, we are able to continue the recursive construction of the desired multiplication. $\square$

Remark 5.1. Using the Φ_h associative multiplications μ_h and $\mu_{t,h}$ from Propositions 5.1 and 5.2 and the equivalence between the monoidal categories $\mathcal{C}(U(\mathfrak{g})[[h]], \Delta, \Phi_h)$ and $\mathcal{C}(U(\mathfrak{g})[[h]], \tilde{\Delta}, \mathbf{1})$ given by the pair (Id, F_h) (see Section 2), one can define $U_h(\mathfrak{g})$ invariant multiplications, either $\mu_h F_h^{-1}$ in the one parameter case or $\mu_{t,h} F_h^{-1}$ in the two parameter case.

References

- [BFFLS] F.Baen, M.Flato, C.Fronsdal, A.Lichnerivicz, D.Sternheimer, Deformation theory and quantization. 1. Deformations of symplectic structures, Ann. Physics 111 (1978), 61-110.
- [Bo] A.Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, Ann. Math., v. 57 (1953), 115-207.
- [Dr1] V.G.Drinfeld, Quasi-Hopf algebras, Leningrad Math. J., 1 (1990), 1419-1457.
- [Do] J.Donin, Double quantization on the coadjoint representation of $sl(n)^*$, Czechoslovak J. of Physics, 47, No 11 (1997), 1115-1122, q-alg/9707031.
- [DG1] J.Donin and D.Gurevich, Quasi-Hopf Algebras and R-Matrix Structure in Line Bundles over Flag Manifolds, Selecta Mathematica Sovietica, 12, No 1 (1993), 37-48.
- [DG2] J.Donin and D.Gurevich, Some Poisson structures associated to Drinfeld-Jimbo R-matrices and their quantization, Israel Math. J., 92, No 1 (1995), 23-32.
- [DGK] J.Donin, D.Gurevich, and S.Khoroshkin, Double quantization of CP^n type orbits by generalized Verma modules, J. of Geom. and Phys., to appear.
- [DGM] J.Donin, D.Gurevich, and S.Majid, R-matrix brackets and their quantization, Ann.de l'Institut d'Henri Poincaré, Phys. Theor. 58, No 2 (1993), 235-246.

- [DGR] J.Donin, D.Gurevich , V.Rubtsov, Quantum hyperboloid and braided modules, in: Algebre non commutative, Groupes Quantiques et Invariants, Societe Mathematique de France, Collection Seminaires et Congres, n 2 (1997), 103-118.
- [DS1] J.Donin and S.Shnider, Quantum symmetric spaces, J. of Pure and Applied Algebra, 100 (1995), 103-115.
- [DS2] J.Donin and S.Shnider, Cohomological construction of quantum groups, Israel Journal of Mathematics, to appear.
- [GP] D.Gurevich and D.Panyushev, On Poisson pairs associated to modified R-matrices, Duke Math. J., v. 73, n. 1 (1994).
- [He] S.Helgason, “Differential geometry, Lie groups, and symmetric spaces”, Academic Press, 1978.
- [KRR] S.Khoroshkin, A.Radul, and V.Rubtsov, A family of Poisson structures on compact Hermitian symmetric spaces, Comm. Math. Phys., v. 152(1993), 299–316.
- [LW] J.H. Lu and A.Weinstein, Poisson-Lie groups, dressing transformations, and Bruhat decompositions, J. of Diff. Geom., 31 (1990), 501-526.
- [NV] O.M.Neroslavsky and A.T.Vlasov, Existence de produits * sur une variété, C.R. Acad. Sci. Paris, 292 I (1981), 71-76.